

Mr Raymond Yap

MBBS, BMedSci, MSurgEd, FRACS, FCSSANZ
Colorectal and General Surgeon
Laparoscopy, Robotics, TEMS/TAMIS, taTME
Open Access Colonoscopy & Gastroscopy
Colorectal Cancer, Diverticulitis, IBD
Haemorrhoids & Anorectal Disease

CR Surgery Clinic

Suite 20, Cabrini Malvern
Isabella Street, MALVERN 3144
Ph: 61 3 8376 6429
Fax: 61 3 9509 0812
Website: <http://crsurgery.com.au>
Email: info@crsurgery.com.au

After Laparoscopy / Laparoscopic Appendicectomy Instructions

You have had your surgery. After the surgery, Mr Yap has talked to your contact person. Any special instructions has been given to you or them. *A responsible adult must take you home. Do not drive or operate machinery for the rest of the day of surgery.*

Diet & Activity

- There are no particular limits on your diet apart from eating healthily
- Avoid activities that put a strain on the abdominal muscles such as sit-ups, weights, and sex. Do not lift anything over 5 kg for 4 weeks. Avoid any activity that causes much pain.
- If you live alone, make arrangements with family or friends in advance to stay and help you.
- Do not drive for the first week, do not go alone the first time and do not go far. Do not drive if you are still on strong painkillers.

Pain and Medications

- **Over the counter pain medications** – Take 1000mg of paracetamol every 6 hours, up to 4 times a day. You can add 400mg of ibuprofen to this, up to 3 times a day – they do not interact.
- **Narcotic pain relief** – If you still have severe pain, use the prescribed tablets every 4-6 hours. *Narcotics can cause constipation but this can be managed by taking laxatives as well.*
- **Stool softeners** – Take a Movicol sachet, once a day if you are on strong painkillers. No script is needed.

Smoking & Alcohol

- **Do not smoke:** If you have been smoking, **stop**. Smoking interferes with healing and increases your risk of complications including infection, hernias, pneumonia, chronic lung disease such as emphysema and lung failure, and cancer. If you need help quitting, talk to your GP.
- **Avoid** alcohol in the first two weeks after surgery. Once you are not taking pain relief, drink no more than one standard alcoholic beverage a day.

Wound Care

- Keep wounds clean and dry. **If there is a dressing please take it off 5 days after your operation.** Leave the wound uncovered unless there is discharge – if so, place plain dry gauze over the wound to absorb this. Do not wear a belt or tight pants against the wound.
- Please bathe everyday. Do not submerge your wounds for 2 weeks. If you have wounds dressings, you can shower and pat dry afterwards.

Things to Watch For

- Heavy wound bleeding/discharge.
- New pain, feeling faint, fevers, or persistent nausea or vomiting.

Follow up

- A follow-up appointment has usually been made for you – see below. If not, please call for a follow up appointment in 4 weeks.
- If you have any questions/concerns, please contact Mr Yap or the rooms.

REVIEW CONSULTATION: Date **TIME:.....**

Suite 20 Cabrini Malvern Isabella St, Malvern / Suite 6 Cabrini Brighton 243 New St, Brighton

Mr Raymond Yap

MBBS, BMedSci, MSurgEd, FRACS, FCSSANZ
Colorectal and General Surgeon
Laparoscopy, Robotics, TEMS/TAMIS, taTME
Open Access Colonoscopy & Gastroscopy
Colorectal Cancer, Diverticulitis, IBD
Haemorrhoids & Anorectal Disease

CR Surgery Clinic

Suite 20, Cabrini Malvern
Isabella Street, MALVERN 3144
Ph: 61 3 8376 6429
Fax: 61 3 9509 0812
Website: <http://crsurgery.com.au>
Email: info@crsurgery.com.au

- If you cannot reach your surgeon or you need immediate attention, please go to your nearest emergency room.

REVIEW CONSULTATION: Date **TIME:.....**

Suite 20 Cabrini Malvern Isabella St, Malvern / Suite 6 Cabrini Brighton 243 New St, Brighton